

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 October 2003 (23.10.2003)

PCT

(10) International Publication Number
WO 03/086471 A2

(51) International Patent Classification⁷: **A61K 47/48**

(21) International Application Number: **PCT/JP03/04745**

(22) International Filing Date: **15 April 2003 (15.04.2003)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
NO. 2002-112864 16 April 2002 (16.04.2002) JP

(71) Applicant (for all designated States except US): **TAN-ABE SEIYAKU CO., LTD.** [JP/JP]; 2-10, Dosho-machi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8505 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ITO, Takahiro** [JP/JP]; 10-1-412, Nishinomiyahama 4-chome, Nishinomiya-shi, Hyogo 662-0934 (JP). **NARISAWA, Shinji** [JP/JP]; 15-103-102, Kamihamuro 3-chome, Takatsuki-shi, Osaka 569-1044 (JP).

(74) Agents: **KAWAMIYA, Osamu** et al.; Aoyama & Partners, IMP Building, 3-7, Shiromi 1-chome, Chuo-ku, Osaka-shi, Osaka 540-0001 (JP).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **LIQUID PREPARATION COMPRISING CAMPTOTHECIN DERIVATIVE AND PHARMACEUTICAL COMPOSITION PRODUCIBLE BY LYOPHILIZING THE PREPARATION**



[I]

(57) Abstract: The present invention relates to a stable liquid preparation comprising a camptothecin derivative which is prepared by binding a compound of the formula [I]; wherein R₁ is a substituted or unsubstituted lower alkyl group, X₁ is a group of the formula: -NHR₂ (R₂ is a hydrogen atom or a lower alkyl group) or a hydroxy group and Alk is a straight or branched chain alkylene group optionally

interrupted by an oxygen atom, and a polysaccharide having carboxyl groups via an amino acid or a peptide, or a pharmaceutically acceptable salt thereof, which is adjusted to pH 5-8, or a stable pharmaceutical composition produced by lyophilizing said liquid preparation.

WO 03/086471 A2


DESCRIPTION

LIQUID PREPARATION COMPRISING CAMPTOTHECIN DERIVATIVE AND
PHARMACEUTICAL COMPOSITION PRODUCIBLE BY LYOPHILIZING THE
5 PREPARATION

TECHNICAL FIELD

The present invention relates to a liquid preparation
comprising a camptothecin derivative or a pharmaceutically
10 acceptable salt thereof, which shows excellent antitumor
activities, a pharmaceutical composition that is producible
by lyophilizing said liquid preparation, and a process for
preparing said pharmaceutical composition.

More particularly, the present invention relates to a
15 liquid preparation for injection comprising a camptothecin
derivative which is prepared by binding a compound of the
formula [I]:



wherein R¹ is a substituted or unsubstituted lower
20 alkyl group, X¹ is a group of the formula: -NHR² (R² is a
hydrogen atom or a lower alkyl group) or a hydroxy group
and Alk is a straight or branched chain alkylene group

optionally interrupted by an oxygen atom,
and a polysaccharide having carboxyl groups via an amino
acid or a peptide, or a pharmaceutically acceptable salt
thereof, which is adjusted to pH 5-8, or a pharmaceutical
5 composition produced by lyophilizing said liquid
preparation, or a process for preparing the same.

BACKGROUND ART

The camptothecin derivatives of the present invention
10 and pharmaceutically acceptable salts thereof are medicinal
substances that show excellent antitumor activities against
various tumors, especially they show excellent therapeutic
effects on solid tumors such as pulmonary cancer, uterine
cancer, ovarian cancer, breast cancer, or gastrointestinal
15 cancer (large bowel cancer, gastric cancer, etc.). It has
been known that said compounds can be administered
parenterally (e.g. intravascular injection) generally in
the form of a liquid preparation (e.g. solution, suspension,
emulsion, etc.) (JP-10-72467A, EP-0757049A).

DISCLOSURE OF INVENTION

The camptothecin derivative above has the structure
wherein a camptothecin compound (active substance) of the
formula [I] is bound to a polysaccharide (carboxymethylated
25 dextran or pullulan) through a spacer (an amino acid or a

peptide). Said camptothecin derivatives, when formulated into a liquid preparation, often undergo hydrolysis at the site of spacer or polysaccharide moiety during the preparation process or storage. Hydrolysis of the polysaccharide moiety results in the reduction of the mean molecular weight of said camptothecin derivatives and the increase of the molecular weight distribution, which variation of molecular weight is apt to affect adversely to the pharmacokinetics of said medicinal substance. Further, hydrolysis of the spacer would result in the release of a considerable amount of an active substance (camptothecin compound [I]) at the time of preparation, which is unfavorable in terms of therapeutic effects or side effects. Accordingly, it was desired to find a liquid preparation that is excellent as to drug stability during the preparation process and storage.

The present inventors have intensively studied to solve the problems above, and have found that a liquid preparation with excellent stability can be obtained by adjusting the pH of a liquid preparation comprising a camptothecin derivative of the present invention between 5 and 8 during the preparation process thereof, and have accomplished the present invention.

That is, the present invention provides a liquid preparation for injection comprising a camptothecin

derivative wherein a camptothecin compound of the formula [I] above is bound to a polysaccharide having carboxyl groups via an amino acid or a peptide, or a pharmaceutically acceptable salt thereof, which preparation is adjusted to pH 5-8.

Further, the present inventors have found that a pharmaceutical composition prepared by lyophilizing the liquid preparation above also shows excellent drug stability during the preparation process and storage. Accordingly, the present invention also provides such a pharmaceutical composition.

MODE FOR CARRYING OUT THE INVENTION

In the present invention, any one(s) of camptothecin derivatives disclosed in JP-10-72467A, that is, camptothecin derivatives wherein a camptothecin compound of the formula [I] above is bound to a polysaccharide having carboxyl groups via an amino acid or a peptide can be used. Specific examples of the camptothecin derivatives include those wherein X^1 of a compound [I] and a carboxyl group of an amino acid or a peptide (e.g. a peptide consisting of 2-5 amino acids) are bound to form an acid-amide bond or an ester bond, and an amino group of said amino acid or peptide and a part or all carboxyl groups of a polysaccharide such as a carboxymethylated dextran or

pullulan are bound to form an acid-amide bond(s).

More specifically camptothecin derivatives include those in which a part or all carboxyl groups of a polysaccharide are bound to a N-terminal amino group of an amino acid or a peptide to form an acid-amide bond, and a C-terminal carboxyl group of said amino acid or peptide is bound with X^1 of a compound of [I] to form an acid-amide bond or an ester bond.

Substituents on a compound of a generic formula [I] include the following substituents. When X^2 is $-NHR^2$, a lower alkyl group in R^2 includes a C_{1-4} alkyl group, and a substituent on a lower alkyl group in R^1 includes a hydroxy group optionally protected, a mercapt group and an amino group (e.g. optionally protected by an alkyl group or an acyl group). Alk includes a straight or branched chain C_{1-6} alkylene group which is optionally interrupted by an oxygen atom.

Polysaccharides related to the present invention include a polysaccharide having originally a carboxyl group in its molecule (e.g. hyaluronic acid, pectin, etc.), a polysaccharide (e.g. carboxymethylated pullulan, carboxymethylated dextran, etc.) which is prepared by introducing a carboxyl group into a polysaccharide having originally no carboxyl group in its molecule (e.g. pullulan, dextran, etc.). Among them carboxymethylated dextran (e.g.

degree of carboxymethylation is more than 0.3 and less than 0.8) is especially preferable. Its mean molecular weight is preferably 20,000 - 400,000, especially preferably 50,000 - 150,000.

5 Preferable camptothecin derivatives are those wherein R^1 is an unsubstituted C_{1-6} alkyl group, X^1 is an amino group and Alk is a straight chain C_{1-6} alkylene group not interrupted by an oxygen atom, a polysaccharide is a carboxymethylated dextran or pullulan, and a peptide is a
10 peptide consisting of 2 - 5 amino acids.

 More preferable camptothecin derivatives are those wherein R^1 is ethyl group, a group of the formula: X^1 -Alk-O- is 3-aminopropoxy group, and camptothecin compound [I] bound at position 10 of a camptothecin nucleus and dextran
15 in which a carboxyl group is introduced, are bound via a peptide selected from a group consisting of glycyl-glycyl-L- or D-phenylalanyl-glycine, glycyl-glycine, glycyl-glycyl-glycine, glycyl-glycyl-glycyl-glycine, glycyl-glycyl-glycyl-glycyl-glycine, L- or D-phenylalanyl-glycine,
20 and L- or D-leucyl-glycine. Among those peptides, glycyl-glycyl-glycine is especially preferable.

 As pharmaceutically acceptable salts of camptothecin derivatives, alkali metal salts such as sodium salt or potassium salt, alkaline earth metal salts such as calcium
25 salt, or amino acid salts such as arginine salt or lysine

salt are illustrated.

The liquid preparation of the present invention is prepared, for example as follows; (1) a camptothecin derivative above or its pharmaceutically acceptable salt and if necessary other ingredients (e.g. excipients for the pharmaceutical preparations such as buffer, a stabilizing agents) are dissolved in a liquid medium such as water for injection etc., (2) the solution is adjusted to pH 5-8, preferably 5-7.5, more preferably 5-7, especially preferably 6-7 with a suitable buffer (e.g. citric acid, hydrochloric acid, sodium hydroxide, etc.), and then, (3) after diluted with water for injection to get desired drug concentration, the solution is filtered through a membrane filter etc., to remove the insoluble materials (pyrogen etc.) and then is filled into a sealing glass vessel, followed by sterilization to prepare the liquid preparation.

The amount of a camptothecin derivative or a pharmaceutically acceptable salt thereof is not limited, but is 1% (w/v) to 20% (w/v), preferably 1% (w/v) to 10% (w/v).

Buffer used for the liquid preparation of the present invention is selected from the group consisting of citric acid, an alkali metal citrate (e.g. sodium citrate etc.), acetic acid, an alkali metal acetate (e.g. sodium acetate etc.), and an alkali metal dihydrogen phosphate (sodium

dihydrogen phosphate etc.). These compounds are suitably combined to use as the buffer. The preferable combination as the buffer is a combination of citric acid and sodium citrate, a combination of citric acid and sodium dihydrogen phosphate, and a combination of acetic acid and sodium acetate, preferably a combination of citric acid and sodium citrate. Ionic strength of the buffer used for the liquid preparation of the present invention can be adjusted to, for example, 0.01-0.6, preferably 0.01-0.3, especially preferably 0.05-0.2.

To the liquid preparation of the present invention and the lyophilized composition thereof can be added conventional ingredients used for injection as well as the above mentioned ingredients. These ingredients are fillers (lactose, sucrose, mannitol, dextran, maltose, trehalose, etc.), solubilizing agents (polyoxyethylene solbitan fatty acid ester such as polysorbate 80 etc., polyoxyethylene hydrogenated castor oil such as HCO-60 etc, polyoxyethylene alkyl ether such as polyoxyethylene lauryl ether, solbitan fatty acid ester such as Span 80 etc.), stabilizer (alkali metal carbonate such as sodium carbonate, alkali hydrogen carbonate such as sodium hydrogen carbonate etc.), antioxidants (cysteine hydrochloride, tocopherol, ascorbic acid, etc.), tonicity agents (glycerin, glucose, etc.), and preservatives (thimerosal, ethanol, propylene glycol,

benzyl alcohol, para hydroxybenzoic acid alkyl ester such as para hydroxybenzoic acid butyl ester, etc.).

The amount of the filler is, for example, 10-100% to a camptothecin derivative [I] or a pharmaceutically acceptable salt thereof. The amount of the solubilizer is, for example, 0.1-10% to a camptothecin derivative [I] or a pharmaceutically acceptable salt thereof. The amount of the stabilizer is, for example, 0.1-10% to a camptothecin derivative [I] or a pharmaceutically acceptable salt thereof. The amount of the antioxidant is, for example, 0.1-10% to a camptothecin derivative [I] or a pharmaceutically acceptable salt thereof. The amount of the tonic agent is for example, 0.01-1% to a camptothecin derivative [I] or a pharmaceutically acceptable salt thereof. The amount of the preservative is, for example, 0.001-0.2% to a camptothecin derivative [I] or a pharmaceutically acceptable salt thereof.

The liquid preparation prepared above is filled into a hard vessel such as a sterile ampoule, a vial, a syringe, etc., and is lyophilized by a conventional method to prepare the pharmaceutical composition of the present invention.

The lyophilized pharmaceutical composition of the present invention is prepared as follows.

The amount of the liquid preparation to be filled into

a vessel is, for example, preferably 5-50%(v/v) per the volume of the vessel, especially preferably 10-25%(v/v).

The external temperature on lyophilization is kept preferably at -50 to 60°C, especially preferably -50 to 40°C, and the pressure for sublimation of the solvent used is preferably 0.01-0.2 Torr, more preferably 0.01-0.1 Torr. The rate of lyophilization is preferably adjusted such that the volume of the solvent (calculated into a solution) is sublimated at the rate of 10 μ l to 100 μ l per 1cm² of the surface area from which the solvent is sublimated for one hour, especially 30 μ l to 60 μ l under controlling ingredients of the liquid to be lyophilized, temperature at lyophilization, pressure at sublimation of the solvent, etc.

In case of lyophilizing the liquid preparation, especially the preparation containing mannitol, dextran, and/or sodium carbonate, etc., the breakage of the vessel is protected by previously adding at least one salt selected from the group consisting of alkali metal chlorides (lithium chloride, sodium chloride, potassium chloride, etc.), alkaline earth metal chlorides (magnesium chloride, calcium chloride, etc.) and alkali metal sulfates (lithium sulfate, potassium sulfate, sodium sulfate, etc.), to said liquid preparation. In this case, preferable salts are sodium chloride, sodium sulfate, etc. The amount of said salt is preferably 0.01-10%, more preferably 0.1-5%

per the drug (weight).

The liquid preparation and the pharmaceutical composition prepared by lyophilizing the liquid preparation are preferably stored in a light resistant sealing vessel.

5 The liquid preparation of the present invention as prepared above, has an excellent property as to drug-stability (a camptothecin derivative) during the preparation process or storage. Therefore, the liquid preparation can be administered directly to a patient. The
10 dosage of the liquid preparation is varied on age, body weight, or condition, but is usually 0.02-50mg, especially 0.1-10mg/kg in calculation to a camptothecin compound [I] (in case of X^1 being $-NHR^2$, its hydrochloride).

 The pharmaceutical composition prepared by
15 lyophilizing the liquid preparation of the present invention, has also an excellent property as to drug-stability during the preparation process or storage, and therefore, it is useful for an injection prepared when necessary.


20 The present invention is further explained in detail by examples, but the present invention should not be limited by these examples.

Example 1

25 Preparation for liquid preparations

Based on ingredients of Table 1 below, an aqueous drug solution was prepared and filtered through a membrane filter (type: GS, pore diameter: 0.22 μ m prepared by Millipore Ltd.). The filtrate (1mL) was filled into a glass 3mL-ampoule. Each ampoule was sterilized in vapor at 100°C for 15 min to obtain a liquid preparation.

Drug: Camptothecin derivative described in Example 84 of Jp-10-72467A as represented by the following formula:



wherein CM means "carboxymethylated".

Table 1

	Comparative example	Liq. preparation of present invention			
		1	2	3	4
Drug (g)		0.4			
Sodium dihydrogen phosphate(g)	0.110	0.147	0.180	0.213	0.245
Citric acid	0.118	0.093	0.071	0.047	0.023
0.4M Aq. sodium dihydrogen phosphate solution	q.s.	q.s.			
0.2M Aq. citric acid solution	q.s.	q.s.			
Sodium chloride(g)	0.771	0.771			
Water for injection	q.s.	q.s.			
Total	100mL	100mL			
pH	4.0	5.0	6.0	7.0	8.0

Stability of liquid preparations

The preparation prepared above was preserved under each preservation condition (at 60°C for 20 days, 50°C for 30 days or 40°C for 120 days), and the stability of the drug was tested (Mean molecular weight and molecular weight distribution, and amount of free active camptothecin). The
5 result was shown in the following Table 2. The mean molecular weight of the drug was calculated by GPC multi angles Laser scattering method (MALLS method) and the mean molecular weight distribution was calculated by the
10 following formula:

Mean molecular weight distribution = weight of mean
molecular weight (MW)/number of mean molecular weight (MN)

Table 2

	pH	Preservation condition	Mean molecular weight	Mean molecular weight distribution
Liq. preparation 1 of present invention	5.0	Initial	138,900	1.195
		60°C for 20 days	125,800	1.183
Liq. preparation 2 of Present invention	6.0	Initial	129,100	1.169
		60°C for 20 days	131,200	1.177
Liq. preparation 3 of Present invention	7.0	Initial	131,400	1.191
		60°C for 20 days	131,400	1.186
Liq. preparation 4 of Present invention	8.0	Initial	130,900	1.202
		60°C for 20 days	127,000	1.195
Comparative example	4.0	Initial	129,800	1.200
		60°C for 20 days	110,100	1.720

Table 3

	pH	Amount of free active camptothecin compound (%) *			
		Initial	60°C for 20 days	50°C for 30 days	40°C for 120 days
Liq. preparation 1 of present invention	5.0	3.08	13.15	9.17	10.60
Liq. preparation 2 of present invention	6.0	1.67	8.12	5.83	5.53
Liq. preparation 3 of present invention	7.0	1.48	14.53	6.60	6.52
Liq. preparation 4 of present invention	8.0	1.93	17.32	8.31	7.95
Comparative example	4.0	13.81	23.73	24.84	27.33

* : Active camptothecin compound means a compound of the following formula and the amount was quantitatively analyzed by the following conditions (the same hereinafter).


Quantitative analysis: A sample solution was diluted with 0.2M formic acid-ammonium formate buffer in 200 times and then, the diluted solution (0.4mL) and an internal standard solution (0.1mL) were mixed and the mixture was filtered through a membrane filter (pore diameter; 0.45 μ m) to prepare a test sample for quantitative analysis. The sample was quantitatively analyzed by subjecting to HPLC under the following conditions.

The amount (%) of free active camptothecin in each sample was calculated as 100% of the amount of free active camptothecin compound produced by adding 10 times amount of

6N hydrochloric acid to the sample solution preserved in a refrigerator and then heating at 100°C for 4 hours.

HPLC conditions:

- Column: Inertsil ODS (prepared by GL Science Inc.)
- Mobile phase: 35mM formic acid-ammonium formate buffer (pH3)/acetonitrile=80/20 (flow: 1.0mL/min.)
- Column temperature: 40°C
- Detection: Fluorophotometer (Ex=360, Em=420nm)
- Active camptothecin compound:



wherein Ra is hydrogen atom, Gly-, Gly-Gly- or Gly-Gly-Gly-.

From the result above, in the liquid preparations of the present invention (pH 5-8), decrease of the mean molecular weight of the drug is less in comparison with a liquid preparation of the comparative example and therefore, increase of the molecular weight distribution of the drug was recognized being protected. This reveals that in the liquid preparation of the present invention degradation of the drug (namely cleavage of chain of dextran molecule) can be prevented and undesired formation of free active camptothecin compound due to degradation of spacer portion can be also prevented.

Example 2

Preparation of lyophilized compositions

Using the same drug as the drug of Example 1, and

based on ingredients described in Table 4 each aqueous drug solution was prepared and filtered through a membrane filter (type: GS, pore diameter: 0.22 μ m prepared by Milipore Ltd.). The filtrate (1mL), was filled into a colorless 13-mL vial and the vial was sealed. Each vial was subjected to lyophilization (pre-freezing: -50°C for 3 hours, primary dehydration: 20°C for 30 hours, secondary dehydration: 60°C for 6 hours) to prepare a lyophilized drug composition.

Table 4

	Comparative example		Composition of present invention			
	A	B	1	2	3	4
Drug (g)	5.0					
Sodium dihydrogen phosphate (g)	0.0059	0.110	0.147	0.180	0.213	0.245
Citric acid	0.153	0.118	0.093	0.071	0.047	0.023
0.4M Aq. Sodium dihydrogen phosphate solution	q.s.		q.s.			
0.2M Citric acid solution	q.s.		q.s.			
Water for injection	q.s.		q.s.			
Total	100mL		100mL			
pH	3.0	4.0	5.0	6.0	7.0	8.0

Stability of lyophilized compositions

The preparations prepared above were preserved at 60°C for 20 days, and the stability of the drug compositions was tested (Change of color, Insoluble materials are present or not after reconstitution, molecular weight distribution of the drug, and amount of free active compound). The result

was shown in the following Tables 5-1 and 5-2.

Table 5-1

[Presence of insoluble materials or not]

	Comparative example		Composition of present invention			
	A	B	1	2	3	4
Change of color	No (yellow)	No (pale yellowish green)	No (pale yellowish green)			
State after reconstitution	Insoluble material: yes	Insoluble material: slight	Insoluble material: no			
pH after reconstitution	3.0	4.1	5.1	6.1	7.1	8.1

5

Table 5-2

[Change of mean molecular weight and molecular weight distribution of drug]

	pH	Condition of preservation	Mean molecular weight	Mean molecular weight distribution
Composition of present invention 1	5.0	Initial	135,400	1.144
		60°C for 20 days	128,700	1.145
Composition of present invention 2	6.0	Initial	132,800	1.145
		60°C for 20 days	128,500	1.140
Composition of present invention 3	7.0	Initial	130,600	1.143
		60°C for 20 days	128,300	1.147
Composition of present invention 4	8.0	Initial	129,800	1.144
		60°C for 20 days	131,100	1.128
Comparative example A	3.0	Initial	132,900	1.120
		60°C for 20 days	150,300	1.280
Comparative example B	4.0	Initial	135,100	1.134
		60°C for 20 days	138,100	1.209

Table 5-3

[Amount of free active compound]

	pH	Amount of free active compound (%) (Preserved conditions: 60°C for 20 days)
Composition 1 of present invention	5.0	<0.3
Composition 2 of present invention	6.0	<0.3
Composition 3 of present invention	7.0	<0.3
Composition 4 of present invention	8.0	<0.3
Comparative example A	3.0	0.76
Comparative example B	4.0	0.48

Example 35 Preparation of lyophilized compositions

The same drug as example 1 (10g), citric acid monohydrate (0.42g), and sodium chloride (500mg) were dissolved in water for injection (100mL) and the solution was adjusted to pH 5.0 with 1M sodium hydroxide to make the total volume 200mL by adding water for injection. The solution was filtered through a membrane filter (type: GS, pore diameter: 0.22 μ m prepared by Milipore Ltd.) and the filtrate (2ml), was filled into a colorless glass 3-mL ampoule. Each ampoule was lyophilized by a usual method to prepare a lyophilized preparations prepared when necessary (the preparation of the present invention).

As a comparative example, the same drug (10g) as used in example 1, and citric acid monohydrate (0.42g) were

dissolved in water for injection (100mL) and the solution was treated by the same manner as mentioned above to prepare lyophilized preparations prepared when necessary (Sodium chloride was not added.).

5 The breakage of the glass ampoules was tested on the composition of the present invention and the composition of the comparative example. The result was shown in the following Table 6.

Table 6

	Broken number per 100 ampoules
Lyophilized composition of the present invention	0
Lyophilized composition of Comparative example	40

Example 4

Preparation of lyophilized compositions

15 The same drug as example 1 (5g), citric acid monohydrate (0.093g), anhydrous sodium dihydrogen phosphate (0.147) and sodium chloride (50mg) are dissolved in water for injection (50mL) and the solution is adjusted to pH 5.0 with 0.4M aqueous sodium dihydrogen phosphate solution or 0.2M aqueous citric acid solution to make the total volume 100mL by adding water for injection. The solution is
20 filtered through a membrane filter (type: GS, pore diameter: 0.22 μ m prepared by Milipore Ltd.) and the filtrate (20ml) is filled into a glass 100-mL vial. Each vial is lyophilized by a usual method to prepare lyophilized compositions prepared when necessary.

Example 5

Preparation of lyophilized compositions


The same drug as example 1 (5g), citric acid monohydrate (0.093g), sucrose (5g) and sodium chloride (50mg) are dissolved in water for injection (50mL) and the solution is adjusted to pH 6.0 with 1M aqueous sodium hydroxide solution to make the total volume 100mL by adding water for injection. The solution is filtered through a membrane filter (type: GS, pore diameter: 0.22 μ m prepared by Milipore Ltd.) and the filtrate (20mL) is filled into a glass 100mL-vial. Each vial is lyophilized by a usual method to prepare lyophilized composition prepared when necessary.

EFFECT OF THE INVENTION

The liquid preparation of the present invention and the composition prepared by its lyophilization have an excellent effect that the degradation of the drug (camptothecin) is less in any stage such as its preparation process, distribution and preservation.

CLAIMS

1. A liquid preparation comprising a camptothecin derivative which is prepared by binding a compound of the formula [I]:



wherein R^1 is a substituted or unsubstituted lower alkyl group, X^1 is a group of the formula: $-NHR^2$ (R^2 is a hydrogen atom or a lower alkyl group) or a hydroxy group and Alk is a straight or branched chain alkylene group optionally interrupted by an oxygen atom, and a polysaccharide having carboxyl groups via an amino acid or a peptide, or a pharmaceutically acceptable salt thereof, which is adjusted to pH 5-8.

2. The liquid preparation according to claim 1 wherein one or more compounds selected from the group consisting of citric acid, an alkali metal citrate, acetic acid, an alkali metal acetate and an alkali metal dihydrogen phosphate are used as the buffer.

3. The liquid preparation according to claim 2 wherein ionic strength of the buffer is 0.2 or less than 0.2.

4. The liquid preparation according to any one of claims 1

to 3 wherein the pH is adjusted to 5 to 7.5.

5. The liquid preparation according to any one of claims 1 to 3 wherein the pH is adjusted to 5 to 7.

6. The liquid preparation according to any of claims 1 to 3 wherein the pH is adjusted to 6 to 7.

7. The liquid preparation according to any one of claims 1 to 6 wherein the amount of the camptothecin derivative or its pharmaceutically acceptable salt is 1% to 20%.

8. The liquid preparation according to any one of claims 1 to 7 wherein one ore more ingredients selected from a stabilizer and a filler are further contained.

9. The liquid preparation according to any one of claims 1 to 8 wherein one ore more stabilizers selected from an alkali metal carbonate and alkali metal hydrogen carbonate, and one ore more fillers selected from lactose, sucrose, mannitol, dextran, maltose and trehalose are further contained.

10. The liquid preparation according to any one of claims 1 to 9 wherein one or more salts selected from an alkali metal chloride, an alkaline earth metal chloride and an alkali metal sulphate are further contained.

11. The liquid preparation according to claim 1 wherein R^1 is an unsubstituted C_{1-6} alkyl group, X^1 is an amino group and Alk is a straight chain C_{1-6} alkylene group not interrupted by an oxygen atom, a polysaccharide is a

carboxymethylated dextran or pullulan, and a peptide is a peptide consisting of 2 - 5 amino acids.

12. The liquid preparation according to claim 11 wherein R¹ is ethyl group, a group of the formula: X¹-Alk-O- is 3-

5 aminopropoxy group, and the camptothecin compound [I] is bound at position 10 of a camptothecin nucleus, the polysaccharide is dextran in which a carboxyl group is introduced, the peptide is glycyl-glycyl-L- or D-phenylalanyl-glycine, glycyl-glycine, glycyl-glycyl-glycine, 10 glycyl-glycyl-glycyl-glycine, glycyl-glycyl-glycyl-glycyl-glycine, L- or D-phenylalanyl-glycine, and L- or D-leucyl-glycine.

13. The liquid preparation according to claim 12 wherein the peptide is glycyl-glycyl-glycine.

15 14. A lyophilized drug composition prepared by lyophilizing the liquid preparation according to any one of claims 1 to 13.

15. A liquid composition for injection wherein the composition according to claim 14 is dissolved in an 20 aqueous medium.

1/2

SEQUENCE LISTING

<110> TANABE SEIYAKU CO., LTD.; ITO, Takahiro; NARISAWA,
Shinji

<120> Liquid preparation comprising camptothecin derivative
and pharmaceutical composition producible by lyophilizing
the preparation

<130> 663732

<150> JP 2002-112864

<151> 2002-04-16

<160> 2

<210> 1

<211> 4

<212> PRT

<213> Artificial sequence

<220>

<223> peptide spacer existing between a polysaccharide
halving a carboxyl group and an active substance having
anti-tumor activity

<400> 1

Gly Gly Gly Gly

1

2/2

<210> 2

<211> 5

<212> PRT

<213> Artificial sequence

<220>

<223> peptide spacer existing between a polysaccharide
having a carboxyl group and an active substance having an
anti-tumor activity

<400> 2

Gly Gly Gly Gly Gly

1

5

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
23 October 2003 (23.10.2003)

PCT

(10) International Publication Number
WO 2003/086471 A3

(51) International Patent Classification⁷: **A61K 47/48**,
47/12, 47/02

(21) International Application Number:
PCT/JP2003/004745

(22) International Filing Date: 15 April 2003 (15.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
NO. 2002-112864 16 April 2002 (16.04.2002) JP

(71) Applicant (for all designated States except US): TAN-
ABE SEIYAKU CO., LTD. [JP/JP]; 2-10, Dosho-machi
3-chome, Chuo-ku, Osaka-shi, Osaka 541-8505 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ITO, Takahiro
[JP/JP]; 10-1-412, Nishinomiyahama 4-chome, Nishi-
nomiya-shi, Hyogo 662-0934 (JP). NARISAWA, Shinji
[JP/JP]; 15-103-102, Kamihamuro 3-chome, Takat-
suki-shi, Osaka 569-1044 (JP).

(74) Agents: KAWAMIYA, Osamu et al.; Aoyama & Partners,
IMP Building, 3-7, Shiromi 1-chome, Chuo-ku, Osaka-shi,
Osaka 540-0001 (JP).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI,
SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report

(88) Date of publication of the international search report:
15 April 2004

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: LYOPHILIZED AND LIQUID PREPARATIONS COMPRISING A POLYSACCHARIDE DERIVATIVE OF CAMP-
TOTHECIN



[I]

(57) Abstract: The present invention relates to a stable liquid preparation comprising a camptothecin derivative which is prepared by binding a compound of the formula [I]: wherein R₁ is a substituted or unsubstituted lower alkyl group, X₁ is a group of the formula: -NHR₂ (R₂ is a hydrogen atom or a lower alkyl group) or a hydroxy group and Alk is a straight or

branched chain alkylene group optionally interrupted by an oxygen atom, and a polysaccharide having carboxyl groups via an amino acid or a peptide, or a pharmaceutically acceptable salt thereof, which is adjusted to pH 5-8, or a stable pharmaceutical composition produced by lyophilizing said liquid preparation.

WO 2003/086471 A3

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/JP 03/04/45

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61K47/48 A61K47/12 A61K47/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 03 015826 A (TANABE SEIYAKU CO) 27 February 2003 (2003-02-27) page 1, line 5 - line 20 page 10, line 1 - line 8 page 14, line 7 - line 14 page 15 page 23 - page 25 claims	1, 11-13
E	EP 1 308 171 A (DAIICHI SEIYAKU CO) 7 May 2003 (2003-05-07) page 2, paragraph 1 - paragraph 10 page 3, paragraph 26 - page 4, paragraph 30 page 5, paragraph 39 - paragraph 42 claims; examples	1-6, 11-15
X	& WO 02 05855 A (DAIICHI PHARMACEUTICAL CO., LTD) 24 January 2002 (2002-01-24) --- -/-	1-6, 11-15

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

- *T* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the International search

7 October 2003

Date of mailing of the International search report

05/11/2003

Name and mailing address of the ISA

European Patent Office, P.O. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Rankin, R

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/JP 03/04745

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	EP 1 310 254 A (DAIICHI SEIYAKU CO) 14 May 2003 (2003-05-14) page 2, paragraph 1 - paragraph 3 page 5, paragraph 42 - paragraph 44 claims; examples	1-15
A	& WO 01 076603 A (DAIICHI PHARMACEUTICAL CO., LTD) 18 October 2001 (2001-10-18) -----	1-15
A	HARADA M ET AL: "Determinants for the drug release from T-0128, camptothecin analogue-carboxymethyl dextran conjugate" JOURNAL OF CONTROLLED RELEASE, ELSEVIER SCIENCE PUBLISHERS B.V. AMSTERDAM, NL, vol. 69, no. 3, 3 December 2000 (2000-12-03), pages 399-412, XP004221290 ISSN: 0168-3659 page 399 -page 412 -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PC1/JP 03/04745

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03015826	A	27-02-2003	WO 03015826 A1	27-02-2003
			JP 2003137818 A	14-05-2003
			US 2003092608 A1	15-05-2003
EP 1308171	A	07-05-2003	AU 7103701 A	30-01-2002
			BR 0112417 A	01-07-2003
			CA 2415922 A1	24-01-2002
			EP 1308171 A1	07-05-2003
			NO 20030139 A	13-03-2003
			WO 0205855 A1	24-01-2002
			US 2003148931 A1	07-08-2003
EP 1310254	A	14-05-2003	AU 4874301 A	23-10-2001
			BR 0109891 A	31-12-2002
			CA 2405453 A1	03-10-2002
			EP 1310254 A1	14-05-2003
			NO 20024823 A	06-12-2002
			CN 1433310 T	30-07-2003
			WO 0176603 A1	18-10-2001
			US 2003109460 A1	12-06-2003